

## BioPro-Bokashi usage guide

### Dosage and Direction for use:

- BioPro-Bokashi is ready to use,
- Mix only as much as is directed in the usage guide for each application

*Note: Mixture should be used within 72 hours*

	Species	Route	Dosage	Directions
Poultry (Broilers)	Hatch to 21 days	Feed	10kg per 1 ton feed	Mix into feed prior to feeding.
	21+ days	Feed	5kg per 1 ton feed	
Poultry (Layers)	Hatch to Pullet Rearing	Feed	10kg per 1 ton feed	Mix into feed prior to feeding.
	Housing to 40 weeks	Feed	10kg per 1 ton feed	
	40 weeks + (During Laying Cycle)	Feed	5kg per 1 ton feed	
	Stress Periods	Feed	15kg per 1 ton feed	



BioPro-Bokashi is a feed additive concentrate of probiotic microorganisms for administration via animal feed.

# BioPro-Bokashi

## Probiotic Feed Additive

Minimum Content Lactic Acid Bacteria:  $3 \times 10^5$  cfu/ml

**BIOPRO-BOKASHI IS NOT HAZARDOUS**

### Storage

Store at room temperature out of direct sunlight. Sedimentation may form. This is normal as this is a natural product.

**Registration holder, manufactured and distributed by: Rhino Agrivantage (Pty) Ltd.**  
**Registration no. 2010/ 005404/ 07**

Tel: +27 21 975 4633

Email: [info@rhinoagrivantage.co.za](mailto:info@rhinoagrivantage.co.za)

Address: 5 Waterblommetjie Street,  
Wellington Industrial Park, Wellington, 7654

Registration No: V27546

Act 36 of 1947.



*Please recycle*

**SHAKE WELL BEFORE USE**

1kg     3kg     5kg     25kg